Objective: To estimate the number of gypsy moth egg masses above

and below snowline

Time of Year: August-April; whenever the ground is snowfree

Equipment Needed: Rope to measure 18.6'

2' measure

Baggie or other container to collect egg masses

Data Sheets Needed: "Gypsy Noth Survey"

Procedure: See "Instructions for Gypsy Moth Survey; September

1978", attached. See "Gypsy Moth Egg Mass Counting" for

information on aging egg masses

Interpreting Results:

Over 500 egg masses/acre has the potential for significant

defoliation

Less than 200 pag mage

Instructions for Gypsy Moth Survey September 1978

	2. Species Composition: Estimate the percentage of major tree species in	
	THE APPENDICTOR OF THE PROPERTY OF THE PROPERT	
, ² (2		
	,	Y
-		Į
·		
- . ,		
-		
<u></u>		
•		
1.		
· ^		
` <u></u>	• • • • • • • • • • • • • • • • • • •	
P-	7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	
(Tax		
Į.		
-)/-		
<u>, E</u>		
<u> </u>		
*		
- 1		
. 1		
i. 1		
- 		
*		
<u></u>		
;		
.		

egg mass plots unless requested by the owner. Do record all other data. If egg mass plots are justified, you may wish to determine species composition after you've established the plots. Species composition does not have to add up to 100%.

3. Plot Data: Randomly establish 1/40 acre plots using the following

	BLOCK NAME BLOCK SIZE(acres) SPECIES COMPOSITION: OAK BIRCH	EVALUATOR(s) ROCK OUTCROPS(circle) present absent BASAL AREA or % OF TOTAL
	ASPEN OTHER MAJOR SPECIES(list) NUMBER OF EGG MASSES PER FIVE MINUTE W LINE 1 LINE 2 LINE 3 LINE LOCATION OF EGG MASSES(circle) Tookle	4 LINE 5 LINE 6 LINE 7 LINE 8
ž.		_
		\/. ₹
,		
	NUMBER OF NEW EGG MASSES PER 1/40 ACRE PLOT# BELOW 2' ABOVE 2' TOTAL 1	PLOT (18.6' radius) PLOT# BELOW 2' ABOVE 2' TOTAL 6
	AVERACE MIMBER OF ECO MACCED DED LONG	9
-		