

know  
associ-  
s work  
ature,  
a com-  
eastern  
corals,  
[ dates  
lat and  
These  
varia-


st by 3  
*zia fer-*  
er, and  
Pacific  
once at  
op. are  
loporid  
7; Pat-  
nigrate  
male-  
e other  
*rpezia*  
by the  
er and  
vitality  
elonga-

t  
e  
c  
i  
l  
e  
l  
t  
t  
c  
t  
e  
c  
e  
s  
e  
l  
c  
e  
s  
i  
s  
t  
b  
f  
c  
b  
t  
r  
s  
v  
A  
i  
l  
c  
a  
(  
v  
E  
c  
a

ried sig-  
dities of  
(%) were  
for reef  
ever, the

N = 869  
Jun Aug 1989

B



D



a females  
*pezia* post-  
rence of  
*Trapezia*

asonally  
; highest  
females  
g eggs,

females  
gs were  
contin-  
. But this  
d during  
e varied  
cept for  
s appear  
ruitment

so varies  
*allina* to  
percent-  
eef loca-  
1 species  
3 A-C. In  
collected  
emale *T.*  
larvae of  
ne reach  
ary. Our  
increases

△  
△  
△  
△



Fig. 3. Size-f  
Dark bars: fe

during August  
as individual  
seasonal cha  
statistically  
hypothesis  
April or Jun  
size in Janu  
a similar pe  
lowest in Au  
Only 1 juve  
cannot mak  
structure of

*Trapezia f*  
nearly every  
sent on onl  
occurrence i  
but is affect  
present on  
only 37% of

The abund  
with coral l  
higher on the  
on the reef  
 $p < 0.05$ ). Ho  
frequency of  
analysis is re

size

niid  
pora  
rites  
cast-  
it is  
col-  
um-

ads.  
- as  
rom  
only  
onal  
cant  
hest  
 $\bar{x} =$   
.pril  
rant  
dry  
ples  
s in  
cted  
ned  
the  
are  
size,  
size  
htly


mm)  
hout  
and  
essa  
and  
y be  
uce,  
lian  
and  
eau-  
ow-  
Top-  
are  
has  
and  
opsis  
tera,

Log number of eggs

Log number of eggs

Fig. + 4

and  
 basi  
*Har*  
 hea  
 occ  
 cor  
*H*  
 cora  
 cora  
 size  
 (Tal  
 can  
 ( $F_{3,1}$   
 cora  
 2.56  
 moc  
 are  
 thar  
 hete  
 reef  
 O  
 prop  
 The  
 Sex  
 Sea  
 and



n-  
he  
d-  
of  
sh  
an  
es  
8).  
is  
le  
ay

is  
tal  
nd  
is  
ess  
he  
lts  
on,  
aii  
re  
tal

als  
le  
us  
ht  
es

g-  
ds  
tal  
is,  
et,  
k-  
he  
st.  
on  
in,  
is  
he  
ist  
u-  
he  
es.  
er  
be

al-  
ze.  
le,  
ce  
of  
in-

Table 2. Significant variables ( $p < .05$ )

Species
<i>Trapezia</i>
<i>Trapezia</i>
<i>Trapezia</i>
<i>Alpheus</i>
<i>Fennera</i>
<i>Harpiliop</i>
<i>Teleophry</i>

dances and species higher on the Gotelli and A continuous and shallow reef flat contribute to the location.

In spite of the coexisting species their population only to *Trapezia lopora*. Table 2 each species.

One notable head size to pool level, the area nates patterns (Abele, 1976b) affects measurement and Abele, 198 much weaker abundance for correlation coefficient. Coral head size rence of *Teleop*



coral  
tribu-

*sillus*  
na 19:

lism'.

Atoll

fracti-  
a 12:

es in  
ligate  
245  
ceans  
re-  
yura,  
n. Ist.

ciated  
*drina*.

rne et  
ool. 8

n. III.  
nd the  
,  
astern

d with  
in the  
240  
n and  
*pezia*  
ceana

pagos  
itions,

erica.  
pedi-

ra and  
h spe-  
ching

ermin-  
acific.

d non-  
mar.

of host  
ologia

arbor-  
11  
of reef  
ns. 10:

nd the  
rv. 11:

c coral

re  
G  
Gotel  
co  
13  
Holth  
(C  
su  
co  
C  
Huite  
tiv  
Knud  
Br  
lo  
McCl  
C  
QL