nature
o ogy o ·o
ARTICLES
P BLI HED: 18 AP IL 2017 OL ME: 1 A ICLE N MBE : 0132

F

N G [1,2,3]*, L B -P [1]*, P L [4]*, M B [1],
N J. G [5] F . M [1]

Understanding the relationship between biodiversity and ecosystem functioning has been a core ecological research topic over the past decades. Although a key hypothesis is that the diversity of functional traits determines ecosystem functioning, we do not know how much trait diversity is needed to maintain multiple ecosystem functions simultaneously (multifunctionality). Here, we uncovered a scaling relationship between the abundance distribution of two key plant functional traits (specific leaf area, maximum plant height) and multifunctionality in 124 dryland plant communities spread over all continents except Antarctica. For each trait, we found a strong empirical relationship between the skewness and the kurtosis of the trait distributions that cannot be explained by chance. This relationship predicted a strikingly high trait diversity within dryland plant communities, which was associated with a local maximization of multifunctionality. Skewness and kurtosis had a much stronger impact on multifunctionality than other important multifunctionality drivers such as species richness and aridity. The scaling relationship identified here quantifies how much trait diversity is required to maximize multifunctionality locally. Trait distributions can be used to predict the functional consequences of biodiversity loss in terrestrial ecosystems.

Global threats to biodiversity have motivated ecologists to better understand the relationship between biodiversity and ecosystem functioning[1] (BEF). Pioneering BEF experiments conducted in grasslands have shown a positive effect of species richness on ecosystem functioning[2,3], which has been later confirmed in many ecosystems[4–7]. However, there is a general consensus that it is not only the number of species *e e* that influences ecosystem functioning, but also the diversity and the abundance of their functional traits within communities[8–10]. As functional traits relate to how species acquire, share and conserve resources[11], they are often invoked to explain how species assemble within communities[12–14] and impact ecosystem functioning[8–10,15].

Higher trait diversity has been hypothesized to enhance ecosystem functioning because co-occurring species with contrasting trait values may exploit different resources, or the same resources on different spatial or temporal scales[2], and then may increase overall resource utilization[16]. This hypothesis has received support from BEF experiments[17] and empirical surveys[10,18,19]. However, most trait-based BEF studies have been conducted on local or regional scales[17,20], limiting our ability to generalize the effect of trait diversity on ecosystem functioning across ecosystems. Also, many studies have focused on single ecosystem functions (for example, productivity[10]), something that may provide a partial measure or even biased estimate of overall ecosystem functioning if trade-offs or synergies among functions are ignored[4,21,22]. Emerging empirical evidence suggests that higher trait diversity is especially important to sustain multiple ecosystem functions simultaneously[19,23] (multifunctionality). However, the extent to which trait diversity matters for multifunctionality remains unknown on a global scale.

Coupling a global survey of multifunctionality conducted in 124

This inequality generates a mathematical constraint triangle on the possible values of skewness and kurtosis that can be observed for a community (Fig. 2; see also Supplementary Note 1 for detailed information). For a given set of distributions, an increase in skewness^2 will generate an increase in K, all other things being equal. In other words, as the trait–abundance distributions become more skewed, they also become more peaked, indicating a decrease in trait diversity[25]. The slope β of the SKR measures the strength of the relationship, that is, the extent to which trait diversity decreases as trait distributions become more skewed. The -intercept α indicates the lowest kurtosis value at skewness = 0, which corresponds to the highest trait diversity predicted by a given SKR. Importantly, the inequality has a lower boundary that sets a limit to the minimal kurtosis value predicted for any degree of skewness, that is, the potential maximum trait diversity ($K = \text{skewness}^2 + 1$; Fig. 2). Although skewness and kurtosis separately provide valuable information on how trait diversity and abundance is distributed within communities, the SKR reveals the extent to which trait diversity is maximized.

Observed trait distributions in global drylands. The observed 124 trait–abundance distributions were highly heterogeneous and deviated from a symmetric normal distribution in most cases (Fig. 3). Consistent with equation (1), we observed a strong positive relationship between skewness and kurtosis of the distributions for both SLA and H (Fig. 3):

$$K_{\text{SLA}} \approx \text{skewness}_{\text{SLA}}^2 + 2 \quad (2)$$

$$K_H \approx \text{skewness}_H^2 + 3 \quad (3)$$

However, within the constraint triangle, there are a variety of possible SKRs that can be generated (Fig. 2). Empirical SKRs must then be compared with the null distribution of relationships that might arise by sampling random values within the constraint triangle. We used a Monte Carlo analysis to test whether the empirical SKRs differed from random expectations. These randomization tests used three null models that imposed increasing constraints on the species and trait pools.

In the first null model, we simulated the assembly of local plant communities from global pools of species and traits. In this null model, the local species number and trait values are randomly assigned to the null communities according to the hypothesis that the local environment may modify both local trait composition and species richness[38]. In the second null model, we fixed species richness at the site level, but allowed traits to vary across communities to decouple the effect of species richness and trait composition on local trait abundance distributions. In the third null model, we additionally constrained the range of trait values observed within community to be fixed. This third randomization procedure tested whether the distribution of trait values between co-occurring species maximized trait diversity locally (see Methods).

The empirical SKRs for SLA and H deviated strongly from the predictions of all three null models (Fig. 3). The slopes β and the -intercepts α were all lower than expected by chance (Fig. 3 right panels; Table 1 left; see detailed results in Supplementary Note 2; Supplementary Fig. 2). Consequently, observed kurtosis values for each of the 124 trait–abundance distributions for both SLA and H were significantly closer than expected by chance to the lower boundary of the mathematical constraint triangle (Table 1 right; see the analysis on the distance to the lower boundary in Methods). In other words, observed kurtosis within communities was minimal after controlling for the degree of skewness in the data. Deviations from null expectations were consistent for all three null models (Table 1), so the results are robust to the details of how species and trait pools were constructed and randomized.

A general assembly rule in global drylands. Our findings have important implications for understanding community assembly in global drylands. The abundance and the diversity of trait values observed within the 124 communities evaluated cannot be observed by chance, and can be summarized by two empirical SKRs, one for H and another for SLA (Figs 2 and 3). Each of these SKRs defined a family of trait–abundance distributions in which the trait distribution of each studied community represents a particular instance of a more general distribution quantified on the biome scale (Fig. 2). These two biome-scale distributions predict that the kurtosis of

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.

Linking a general assembly rule to ecosystem multifunctionality. We conducted a sensitivity analysis of the best-fitting model to explore how coordinated changes in the skewness and the kurtosis of the trait–abundance distributions affected multifunctionality (model 4, Fig. 4; see Supplementary Note 4 for more details on the sensitivity analysis). The two empirical SKRs predict a peak of multifunctionality to occur at

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.

of xeric shrublands and dry forests (blue and purple diamonds in Fig. 5b, respectively; see also Supplementary Note 4). We observed (1) a maximized multifunctionality state closer to a normal distribution (-intercept $\alpha \approx 3$, along red line in Fig. 5b); and (2) a reduced multifunctionality state closer to the lower boundary (see the shaded area in Fig. 5b). The two empirical SKRs not only show that high trait diversity is needed to sustain multifunctionality, but also precisely quantify how much diversity is required to maximize it locally.

Our study suggests that SLA and *H*—two key plant attributes[26]—can be used as efficient functional markers (*e* , ref. [15]) to scale-up trait diversity to the ecosystem level. b8(o k)(u)0 0 251 T22.3251 w

Using the same randomization test applied across communities (see Methods; Supplementary Note 5), we found that the relationship between trait–abundance distributions observed within communities and multifunctionality also cannot be explained by chance (Fig. 4b). The two empirical SKRs corresponded to the unique sets of trait distributions able to explain multifunctionality with such a high predictive power (Fig. 4b). Results were robust to the three randomization scenarios employed, including the most highly constrained null model based on the local species composition and trait pool observed within communities (Supplementary Note 3). Our analysis thus supports the existence of a unique scaling relationship linking plant functional traits (SLA and *H*), the distribution of their abundance within communities and multiple ecosystem functions.

It is noteworthy that increasing trait diversity to its highest potential values defined by the lower boundary of the constraint triangle does not necessarily increase multifunctionality (Fig. 5b). For example, decreasing the kurtosis value for *H* beyond the normal distribution towards a kurtosis value of 1 (that is, the lower boundary) had a strong detrimental effect on the multifunctionality

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.

We calculated the mean, variance, skewness and kurtosis of the 124 trait–abundance distributions[25,32] for SLA and *H* separately as follows:

$$\text{Mean}_j = \sum_i {}_i T_i \tag{4}$$

$$\text{Variance}_j = \sum_i {}_i (T_i - \text{Mean}_j)^2 \tag{5}$$

$$\text{Skewness}_j = \sum_i \frac{{}_i (T_i - \text{Mean}_j)^3}{\text{Variance}_j{}^{3/2}} \tag{6}$$

$$\text{Kurtosis}_j = \sum_i \frac{{}_i (T_i - \text{Mean}_j)^4}{\text{Variance}_j{}^{2}} \tag{7}$$

where ${}_i$ and T_i

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.

$P(\beta \cap \alpha)$ represents the probability to observe a whole model with both a slope and a -intercept (β_{obs} and α_{obs}) being lower than expected by chance.

Di a ce he e b da : we used a complementary approach to assess whether the observed SKR differs from SKR_{random} by testing whether observed kurtoses of each of the 124 communities sampled were significantly closer to the lower boundary than the random data. The lower boundary defines the lowest

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.

32. Le Bagousse-Pinguet, Y. *e a*. Testing the environmental ltering concept in global drylands.

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved.